

**Literaturverzeichnis zum Artikel „Schröpfen“ auf Carstens-Stiftung.de unter:**

<http://www.carstens-stiftung.de/artikel/schroepfen.html>

Abele J (2003): Das Schröpfen: eine bewährte alternative Heilmethode. 5. Auflage. München: Urban und Fischer.

Bachmann G, Pecker F (1978): Die Schröpfkopfbehandlung. 3. Auflage. Heidelberg: Haug.

Binder A (2008): Neck Pain Clin Evid 08: 1103.

Chirali I (2008): Schröpftherapie in der Chinesischen Medizin. 2. Auflage. München: Urban & Fischer

Côté P, van der Velde G, Cassidy JD et al. (2008): The burden and determinants of neck pain in workers: Results of the Bone and Joint Decade 2000–2010 Task Force on Neck Pain and Its Associated Disorders. Spine 33: 60–S74.

Dobias D (2003): Pneumatische Pulsationstherapie. Teil eines multimodalen Behandlungskonzeptes bei Schmerzen. Praxis Magazin 11: 6–11.

Dworkin RH, Turk DC, Wyrwich KW et al (2008).: Interpreting the clinical importance of treatment outcomes in chronic pain clinical trials: IMMPACT recommendations. J Pain.9: 105–121.

Farhadi K et al. (2009): The effectiveness of wet-cupping for nonspecific low back pain in Iran: A randomized controlled trial. Complementary Therapies in Medicine 17: 9–15.

Fejer R, Kyvik KO, Hartvigsen J (2006): The prevalence of neck pain in the world population: a systematic critical review of the literature. Eur spine J 15: 834–848.

Johnston V, Jimmieson NL, Jull G, Souvlis T (2008): Quantitative sensory measures distinguish office workers with varying levels of neck pain and disability. Pain 137:257–265.

Kim J.I.; Lee M.S.; Lee D.H.; Boddy K.; Ernst E (2009): Cupping for Treating Pain: A Systematic Review. Evid Based Complement Alternat Med. 2009 doi:10.1093/ecam/nep035.

La Touche R, Fernández-de-Las-Peñas C, Fernández-Carnero J, Díaz-Parreño S, Paris-Alemany A, Arendt-Nielsen L (2010): Bilateral mechanical-pain sensitivity over the trigeminal region in patients with chronic mechanical neck pain. J Pain 11: 256–263.

Langevin HM, Stevens-Tuttle D, Fox JR, Badger GJ, Bouffard NA, Krag MH, Wu J, Henry SM (2009): Ultrasound evidence of altered lumbar connective tissue structure in human subjects with chronic low back pain. BMC Musculoskelet Disord. 10, 151–159.

Linton SJ (2000): A review of psychological risk factors in back and neck pain. Spine 25: 1148–1156.

Lüdtke R, Albrecht U, Stange R, Uehleke B (2006): Brachialgia paraesthesia nocturna can be relieved by “wet cupping” – results of a randomised pilot study. Complement Therapies in Medicine 14, 247–253.

MacDermid JC, Walton DM, Avery S, Blanchard A, Etruw E, McAlpine C, Goldsmith CH (2009): Measurement properties of the neck disability index: a systematic review. J Orthop Sports Phys Ther 39: 400–417.

Michalsen A, Bock S, Lüdtke R, Rampp T, Baecker M, Bachmann J, Langhorst J, Musial F, Dobos GJ (2009): Effects of traditional cupping therapy in patients with carpal tunnel syndrome: a randomized controlled trial. J Pain 10: 601–608.

Musial F, Michalsen A, Dobos G (2008): Functional chronic pain syndromes and naturopathic treatments: neurobiological foundations. Forsch Komplementärmed 15:97–103.

Rolke R, Baron R, Maier C, et al (2006): Quantitative sensory testing in the German Research Network on Neuropathic Pain (DFNS): standardized protocol and reference values. *Pain* 123: 231–243.

Schofferman JA, Koestler ME (2005): Whiplash and neck pain-related disability. In: Schultz IZ, Gatchel RJ, editors: *Handbook of complex occupational disability claims: early risk identification, intervention, and prevention*. New York: Springer 203–216.

Scott D, Jull G, Sterling M (2005): Widespread sensory hypersensitivity is a feature of chronic whiplash-associated disorder but not chronic idiopathic neck pain. *Clin J Pain* 21: 175–181.

Woolf CJ, Salter MW (2000): Neuronal plasticity: increasing the gain in pain. *Science* 288: 1765–1769.